

TOBACCO USE

SMOKELESS TOBACCO USE

Erie County Adults Age 18+ 1999, 2001, 2004, 2007

Currently uses smokeless tobacco such as chewing tobacco or snuff

Population	1999		2001		Population	2004		2007	
	%	CI	%	CI		%	CI	%	CI
All Adults	4	3-5	5	4-6	All Adults	2	1-3	3	2-6
Gender					Gender				
Male	8	6-11	9	6-11	Male	4	3-6	7	4-11
Female	0	0-1	1	0-2	Female	0	0-1	0	0-2
Age					Age				
18-29	7	3-11	7	3-11	18-29	3	0-5	n/a	
30-44	7	4-11	8	5-11	30-44	3	1-5	6	3-11
45-64	2	1-3	2	1-4	45-64	2	1-3	2	1-5
65+	3	1-5	2	0-4	65+	1	0-2	1	0-4
Education					Education				
<High School	5	1-9	3	0-6	<High School	1	0-3	n/a	
High School	4	3-6	4	2-6	High School	3	1-5	5	3-10
Some College	3	1-5	5	2-7	Some College	2	0-3	2	0-5
College Degree	6	3-9	7	4-10	College Degree	2	0-3	1	1-4
Income					Income				
<\$10,000	4	0-9	6	0-14	<\$25,000	2	0-3	3	1-11
\$10,000-\$19,999	4	1-7	4	0-8	\$25,000-\$49,999	2	1-4	2	1-5
\$20,000-\$34,999	5	2-7	4	2-7	\$50,000+	3	1-5	5	2-9
\$35,000-\$49,999	5	2-8	6	3-9					
\$50,000+	5	2-8	6	3-9					
Race					Race				
White,non-Hispanic	5	3-6	5	4-6	White,non-Hispanic	2	1-3	3	2-6
Black,non-Hispanic	0		2	0-5	Black,non-Hispanic	0		n/a	
Hispanic	n/a		n/a		Hispanic	^		n/a	

Red indicates significant difference compared to previous year; 95% Confidence Interval (CI)

^Data statistically unreliable; Data excludes missing, don't know, and refused

Source: 1999, 2001, and 2004 Erie County Behavioral Risk Factor Surveillance Survey; 2007 Behavioral Health Risks of Erie County Adults

